

# Administering antibiotics with dry feed

Group-treatment can be administered via the pigs' feed. Do not mix antibiotics in the feed mixer or feeding system.

## The amount of antibiotics required for treatment depends on

- Number of pigs per feeder
- Weight of the pigs
- Kg dry feed consumed per day/feeder

## Mixing antibiotics with dry feed

- Use a cement mixer or a feed cart
- Cement mixer and feed cart must be clearly labelled "Only feed with antibiotics"
- Mix thoroughly to obtain a uniform mixing of feed and antibiotics

## Treatment with antibiotics in dry feed

- Feed with antibiotics is released only in the feeders of the pens with pigs to be treated
- Feed with antibiotics can be fed on solid floor

## Personal safety

- There is a risk of allergic reactions in case of contact with or inhalation of antibiotics
- Wear gloves and mask when mixing antibiotics in feed
- It is recommended to wear a mask when using feed containing antibiotics
- Limit dust production during mixing of antibiotics in feed – use a cover on the cement mixer

## Cross-compliance requirements for treatment with antibiotics in feed:


- Do **not** mix antibiotics in the feed mixer
- Do **not** transport feed with antibiotics in the feeding system
- Comply with the slaughter deadline after antibiotic treatment



Wear mask and gloves when weighing out dry antibiotics



Mix antibiotics and feed very thoroughly



Feed cart labelled "Only feed with antibiotics"



Only give feed with antibiotics to the pigs requiring treatment